

The added value of exhaled breath temperature in respiratory medicineTodor A. Popov¹, Tanya Z. Kralimarkova¹, Marina Labor^{2,3}, Davor Plavec^{3,4}¹Clinic of Allergy and Asthma, Medical University Sofia, Bulgaria²Pulmonology Department, University Hospital Center Osijek, Croatia³Faculty of Medicine, J.J. Strossmayer University, Osijek, Croatia⁴Research Department, Children's Hospital Srebrnjak, Zagreb, Croatia**Abstract**

Recognition of the huge economic burden chronic respiratory diseases pose for society motivated fundamental and clinical research leading to insight into the role of airway inflammation in various disease entities and their phenotypes. However, no easy, cheap and patient-friendly methods to assess it have found a place in routine clinical practice. Measurement of exhaled breath temperature (EBT) has been suggested as a non-invasive method to detect inflammatory processes in the airways as a result of increased blood flow within the airway walls. As EBT values are within a narrow range, the thermometers designed for the purpose of assessing it need to be precise and very sensitive. EBT increases linearly over the pediatric age range and seems to be influenced by gender, but not by height and body weight. In non-smoking individuals with no history of respiratory disease EBT has a natural circadian peak about noon and increases with food intake and physical exercise. When interpreting EBT in subjects with alleged airway pathology, the possibilities of tissue destruction (chronic obstructive pulmonary disease, cystic fibrosis) or excessive bronchial obstruction and air trapping (severe asthma) need to be considered, as these conditions drive (force) EBT down. A prominent advantage of the method is to assess EBT when patients are in a steady state of their disease and to use this "personal best" to monitor them and guide their treatment. Individual devices outfitted with microprocessors and memory have been created, which can be used for personalized monitoring and disease management by telemedicine.

Index Terms: body temperature, thermometry, exhaled breath temperature, airway inflammation, airway remodeling, daily monitoring, personalized medicine, telemedicine.

Introduction

Body temperature: the revival of a hot topic.

Some 200 years ago, when the less numerous human population was plagued by infectious diseases, special focus of the art of medicine involved different patterns of increased body temperature and fever, as evidenced by the monograph of Alexander Philips Wilson, which had multiple editions around the turn of the 18th century [1] (Fig. 1).




Figure 1. A.P. Wilson's "A treatise on Febrile Diseases", 1800

Since that time the control of communicable diseases has dramatically improved, but body temperature measurement has remained an integral part of routine patient examination in clinical practice. The value of 37°C is postulated to be the cut-off point between health and pathology. However, even this most conservative physiological parameter should be perceived as having individual variability, which has to be taken into account when making a judgment about measurements in the febrile range [2].

Humans are warm-blooded species (in scientific terminology "endothermic homeotherms"). Mammalian organisms need to maintain the temperature of vital organs (core body temperature) within a narrow range in order to allow essential enzymatic reactions to occur. The body heat produced as a result of metabolic processes becomes part of a thermal balance, which is regulated through radiation, conduction, convection and/or evaporation [3].

Exhaled breath temperature in humans

1
2
3 The blood circulation and the border surfaces with the ambient environment play a crucial
4
5 role in this heat exchange. Extreme exogenous and endogenous noxious influences can drive
6
7 the core body temperature beyond the narrow range of 33.2–38.2°C [4] and can have fatal
8
9 consequences, while modern day medicine artificially induces hyperthermia in cancer
10
11 treatment [5] or hypothermia to allow prolonged surgical interventions of vital organs [6].
12
13

14
15 For practical purposes thermometry at traditional body sites is performed giving estimates of
16
17 the actual core temperature with a reasonable degree of approximation [4]. Thus, rectal temperature is
18
19 considered most representative of the core body temperature, but its measurement is uncomfortable for
20
21 patients and carries the risk of bacterial contamination. Oral temperature is generally 0.5°C lower than
22
23 the rectal temperature and is more prone to influences from the ambient environment, while the
24
25 temperature of the tympanic membrane taken by infrared tympanic thermometers was found to be
26
27 imprecise compared to rectal values [7].
28
29

30
31 All other conditions of measurement being equal, the differences between the
32
33 temperature values at specific body sites are due to the influence of the “core-to-surface”
34
35 interface. While this may be considered a confounding factor from the viewpoint of
36
37 evaluation of the “true” core temperature, differences due to the core-to-surface gradient may
38
39 present an opportunity to obtain useful information about pathology associated with the
40
41 interface itself. This is particularly relevant for complex anatomical structures, such as the
42
43 lungs, involving vascularized tissues and a plethora of airways of different sizes with
44
45 hierarchical architectonics.
46
47
48
49

The benefits of a ‘breath thermometer’.

50
51
52
53 Perception of the airways as an interface between body core temperature and the
54
55 ambient environment became the rationale for attempts to measure exhaled breath
56
57 temperature (EBT). The deep structures of the lung typically have temperature representative
58
59 of the body core. Its level is determined by the blood flowing along the rich vascular network
60

Exhaled breath temperature in humans

1
2
3 of the alveoli. During breathing, gases and thermal energy are being exchanged between the
4
5 inner milieu of the organism and the ambient environment. The temperature of the inhaled air
6
7 is tempered during its flow in and out of the branching airways, which have a separate system
8
9 of blood supply from the left side of the heart. As blood is the main carrier of thermal energy,
10
11 processes that would modify its flow within the airway walls might reflect on the temperature
12
13 of the outgoing air, i.e. EBT. High precision gauging devices may pick up this signal and
14
15 indicate clinical inferences.
16
17
18
19

Technical aspects of EBT measurement.

20
21
22
23
24 The first experiments assessing EBT were made in conjunction with eNO
25
26 measurement and were conducted in adults by Paredi et al. [8] and in children by Piacentini et
27
28 al. [9]. Both teams used fast reacting thermal sensors placed in front of the mouth of the tested
29
30 subjects, which recorded the rise of EBT during single breath maneuvers and used the mean
31
32 of three exhalation attempts. This required constant temperature of the indoor environment,
33
34 minimal air movement and subject training. Whilst the Paredi team considered the rate of
35
36 increase of EBT as indicative of asthma, the researchers in Verona carried out a series of
37
38 experiments demonstrating that the plateau of the exhaled temperature curve was the variable,
39
40 distinguishing asthmatics from healthy controls [10].
41
42
43
44

45 An alternative approach for EBT measurement was introduced by Popov et al., who
46
47 made use of a specifically designed portable device [11]. It dwelt upon the notion of
48
49 accumulation of the expired thermal energy of the tested subject into an insulated vessel
50
51 containing a heat sink with high thermal capacity, thus making the measurement less
52
53 dependent on ambient factors. The subjects exhaled continuously into the thermal chamber of
54
55 the device until the temperature of the heat sink reached a plateau, indicating that a thermal
56
57 equilibrium was reached inside the closed system. Because of the easy use and acceptability
58
59 by the patients, the instrument allowed repeated measurements over time, with the potential
60

1
2
3 of use as an individual device for measurements at home or in the working environment. Its
4
5 usability was further improved by upgrading the overall design, introducing electronic
6
7 processor and memory allowing automatic detection of the end of measurement, follow up
8
9 and analysis of the temperature curve on the monitor of a computer [12]. Despite these
10
11 technical improvements, the acquisition of measurement skills with the device, by both
12
13 patients and medical personnel, is essential for obtaining meaningful results. A detailed
14
15 description of the engineering aspects of the device and method for EBT measurement has
16
17 been the subject of a specialized review [13]. Further improvements of its applicability
18
19 involved shortening of the time for measurement and rendering it less dependent on the
20
21 conscientious cooperation of the subjects (this latter line of improvement would allow using
22
23 the device in early infancy).
24
25
26
27
28
29

Impact of ambient air temperature on EBT measurement results

30
31
32
33 Even with this closed-circuit multi-breath technique, the issue of the confounding
34
35 influence of the characteristics of the ambient air on the end results of EBT measurement
36
37 remains. As can be expected, the temperature of the ambient air inhaled during the
38
39 measurement affects the accuracy^{1*}, but not the precision^{**} of the measurements. The initial
40
41 analysis of 132 measurements made with EBT as dependent variable and room temperature
42
43 (values on separate days in the range 18–25°C), atmospheric pressure (range 954–982 mbar)
44
45 and humidity (ranges 22–72%) as independent variables, did not detect any of the ambient
46
47 conditions as significant determinants, hence we recommend that the measurements are made
48
49 in a controlled/indoor environment with air temperatures within this range (11).
50
51
52
53
54

55 In a recent real life study Carpagnano et al. measured EBT of 867 volunteers in 3
56
57 different outdoor and indoor (hospital and shopping mall) environments with ambient air
58
59
60

* Accuracy = the degree of closeness of measurements of a quantity to that quantity's true value.

** Precision = the degree to which repeated measurements under unchanged conditions show the same results.

Exhaled breath temperature in humans

1
2
3 temperature ranging from 0 to 38 °C [14]. Out of this random cohort, 298 subjects had never
4
5 smoked and were free of respiratory and other diseases. The regression model with their EBT
6
7 values as a dependent variable and the ambient air temperature as an independent one outlined
8
9 an association, in which the increase of external temperature by 1°C corresponded on average
10
11 to EBT increase of 0.19°C. For this reason it is important to measure the external temperature
12
13 and if necessary to apply a correction factor to the results obtained. Independent technical
14
15 experiments conducted by the manufacturer of X-halo arrived at the same value of this
16
17 conversion coefficient. More data are needed to verify whether this relationship is strictly
18
19 straight linear, or whether an intermediate plateau could exist at room temperature as
20
21 suggested in earlier studies.
22
23
24
25

26
27 Logie et al. demonstrated that the temperature of the inspired air affects both the slope
28
29 and the plateau of EBT and is a significant predictor of EBT in children [15]. This was
30
31 corroborated in the elderly (60-80 years of age) in a study by Bijmens et al. [16].
32
33

34
35 As for atmospheric pressure and humidity, there are no new data to suggest that they
36
37 significantly affect EBT measurement if within a reasonably acceptable range.
38
39

EBT and temperature taken at traditional measurement sites

40
41
42
43 The initial proof-of-concept studies started a long and continuous process of
44
45 assessment of the precision and repeatability of EBT measurements. It was demonstrated that
46
47 the day-to-day measurements in healthy subjects were repeatable with an intraclass
48
49 correlation coefficient of 0.99 [11]. One of the crucial questions, which needed to be
50
51 answered, was whether EBT is just another surrogate measure of core body temperature, or
52
53 whether it also captures the signal emitted by the airways. The pooled analysis of numerous
54
55 EBT and body temperature measurements of healthy subjects and asthmatics did not disclose
56
57
58
59
60

any meaningful correlation between EBT and any of them, while there was a highly significant correlation between otic and axillary temperatures (Table 1).

		Exhaled Breath Temperature
	Axillary Temperature	R=0.01 (P>0.1)
Otic Temperature	R=0.71 (P<0.01)	R=0.06 (P>0.1)

Table 1. The lack of correlation between EBT and body temperatures measured at traditional sites suggests that it is a different physiological indicator.

Partially different results were found by Flouris and Cheung in an experiment with healthy volunteers showing that there was a significant correlation (R=0.58) between rectal temperature and EBT, but changes in EBT were on a larger scale (3 times the change in rectal temperature), thus regulating the core temperature as a reaction to thermal stress [17].

Thus, while core body temperature determines the operative thermal state of humans as a species, EBT represents organ specific physiological modulation of its values, but also reflects pathological changes of the respiratory system.

Physiological factors affecting EBT measurement

Similar to all other new methods with potential clinical applications, EBT measurement requires careful assessment of possible confounding factors to be taken into consideration. EBT is affected by the ambient environment and by different activities in both health and disease states. In an initial report multiple regression analysis did not indicate a significant association of EBT with gender, height, weight, heart rate, blood pressure [11]. Age was a special focus of attention, as the method holds high promise for use in the pediatric population: a positive correlation (R=0.75, P<0.001) was established in healthy children in the age range between 3 and 17 years [18], supported also by the results of Barreto et al. [19] and

Exhaled breath temperature in humans

1
2
3 Vermeulen et al. [20]. In the study of Logie et al., multiple regression analysis indicated slow
4 vital capacity (strongly correlated to age) as a predictor of EBT in a study of 60 children aged
5
6
7
8 between 9 and 11 years of age [15]. Age did not seem to be a major determining factor in the
9
10 pooled analysis of our adult control subjects free of respiratory diseases, but there were
11
12 indications that in elderly people EBT may tend to be lower [16], probably also in relation to
13
14 accompanying geriatric morbidities.
15
16

17
18 Gender is another important determinant of EBT. The pooled analysis we did on all
19
20 our adult healthy control subjects outlined a trend towards somewhat higher EBT in 83 male
21
22 subjects compared with the EBT of 107 women, but it was not statistically significant. In the
23
24 larger study of Carpagnano et al. (143 men and 155 women) a significant gender difference
25
26 emerged with EBT of the male subjects being about 1°C higher [14]. The same was found in
27
28 elderly subjects over 60 years of age [16]. This gender difference may be still preserved in
29
30 asthma according to a cross sectional study involving 69 subjects on maintenance treatment,
31
32 where men had significantly higher EBT [21 Crespo].
33
34
35

36
37 Healthy subjects have different circadian course of EBT compared with their axillary
38
39 temperature: the acrophase (peak temperature) was registered at 19h for EBT and at 13h for
40
41 axillary temperature [22]. The bathyphase (trough temperature) was the same for both
42
43 circadian rhythms at 1h. Repeated measures analysis found both circadian fluctuations to be
44
45 statistically significant. Whether this is also true for patients with inflammatory airway
46
47 disease, remains to be determined.
48
49

50
51 Food intake, especially highly caloric fast utilized carbohydrate products, increase
52
53 EBT within the next hour [23]. Doubling the amount of energy food proportionally increased
54
55 EBT.
56

57
58 Tufvesson et al. found that EBT correlated with an increase in the numbers of club
59
60 cell (Clara) protein (CC16) in plasma and urine after exercise challenge in asthmatics and

Exhaled breath temperature in humans

1
2
3 healthy controls [24]. As CC16 levels in plasma reflect an overall epithelial involvement and
4
5 as no difference between asthmatics and healthy controls appeared, this finding was
6
7 concluded to be a physiological rather than a pathophysiological response.
8
9

10
11 Air pollution from traffic was found to significantly influence EBT in elderly subjects,
12
13 proportionally to the density of the traffic [16]. One other environmental factor affecting the
14
15 respiratory system is tobacco smoke. Smoking the first cigarette for the day was found to
16
17 trigger inflammatory events within the next hour, as evidenced by increase of EBT [25,26].
18
19 Apart from the immediate effect of smoking a cigarette, the issue of the association between
20
21 EBT and cigarette smoking seems to have long term consequences. There was a significant
22
23 inverse correlation between EBT and the number of pack-years in 80 current smokers [25].
24
25 Multiple regression analysis with 'EBT' as dependent variable and 'age', 'gender', 'height',
26
27 'weight' and 'pack-years' as independent variables, identified only 'pack-years' as
28
29 significantly contributing to the overall equation. Similarly, a study replicated these results
30
31 confirming that EBT is sensitive to the acute effect of cigarette smoke, but also found
32
33 significantly higher EBT in current smokers compared to non-smokers and demonstrated that
34
35 after cessation of smoking EBT progressively decreased over time since the last cigarette was
36
37 smoked [27]. The results of a prospective, observational, non-interventional cohort study of
38
39 146 patients, smokers and ex-smokers with a smoking history of >20 pack years without
40
41 chronic obstructive pulmonary disease (COPD) at the start of the study, indicated the
42
43 possibility that the acute effect of smoking one cigarette at baseline can identify the subjects
44
45 who will develop COPD in the course of the two year follow up [28].
46
47
48
49
50
51
52
53

54 In-season high pollen counts increase EBT in sensitized subjects with allergic
55
56 rhinoconjunctivitis with or without asthma [29]. Any natural components of the ambient air
57
58 or gases, aerosolized fluids or particulate matter that can be inhaled accidentally or
59
60 intentionally can potentially influence EBT and need to be specifically explored.

Exhaled breath temperature in humans

1
2
3 Inhalatory therapeutic and diagnostic agents have the potential of impacting EBT due
4 to their effect on bronchial vasculature and airway geometry, which has to be taken into
5 consideration if this method is used for diagnostic and monitoring purposes. In asthmatics
6 and healthy controls inhalation of 400 mcg of salbutamol did not consistently change EBT,
7 about half of the studied asthmatics increased or decreased their EBT beyond the margin of
8 repeatability of this measurement, which was calculated to be $\pm 0.25^{\circ}\text{C}$; whether they
9 represent phenotypes with specific clinical implications remains to be investigated [30]. On
10 the other hand Svensson et al. [31] found that EBT increased after eucapnic voluntary
11 hyperventilation and methacholine challenge test in both asthmatics and healthy subjects
12 representing in their opinion a physiologic vascular effect present after these challenges in the
13 whole respiratory system.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 Body height and body mass index (BMI) do not seem to be significant determinants in
30 both adults and children. An exception is the study of the elderly by Bijmens et al. [16] that
31 pointed out BMI as the EBT predictor, interpreted by the authors as being associated with
32 systemic inflammation. However, the associations between age, gender, weight, height, other
33 physiological indices and EBT need to be revisited with the increasing number of subjects.
34
35
36
37
38
39
40
41

Reference values for EBT in healthy subjects

42 The usability of random single point EBT measurements as an objective marker for
43 diagnostic purposes in respiratory medicine requires establishment of normal reference values.
44 Bearing in mind the multiplicity of the already discussed potential confounders of technical,
45 environmental and physiological nature, this task necessitates sieving through a significant
46 random sample of individuals free of respiratory disease from the general population.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

Detection) and came up with a reference value of EBT in healthy Caucasian non-smoking subjects of $30.459 \pm 2.955^\circ\text{C}$.

Determinants of EBT in respiratory diseases

The airways and deep structures of the lung modulate the gas content and the temperature of the air we inhale so as to ensure optimal gas exchange and to prevent damage to the tissues involved in the process. The mechanisms by which the temperature of the lung tissues is regulated involve modulation of the caliber of the airways and the blood flow through the vast vascular network of the bronchial tree. Pathological processes which affect this intricate regulation would reflect on the overall heat exchange during breathing, turning EBT either up or down (Fig. 3).




Figure 3. EBT determining vectors acting in opposite directions.

‘Turning EBT up’: the inflammatory vector

Vasodilatation is an inherent feature of inflammation, which is a prominent characteristic of asthma and obstructive lung diseases in general [32]. The increased vascularity of the airways in asthma [33] is partly due to the elevated number of vessels

Exhaled breath temperature in humans

1
2
3 associated with angiogenesis and vasodilation caused by the release of mediators, such as
4
5 histamine, bradykinin [34], and nitric oxide (NO) [35]. Acetylcholine is the most important
6
7 mediator to trigger active vasodilation to body heating, although co-transmitters appear to be
8
9 principally involved in the overall response. Vasoactive intestinal peptide, substance P,
10
11 histamine, prostaglandins, and transient receptor potential (TRP) V1 receptor activation seem
12
13 to be included. There appears to be a role for nitric oxide in active vasodilation, as the
14
15 response is attenuated by nitric oxide synthase inhibition [36].
16
17
18
19

20 The relationship between the level of EBT and the bronchial blood flow, presumably
21
22 due to increased vascularity, has been clearly demonstrated in a clinical experiment [37]. The
23
24 level of exhaled nitric oxide (eNO) was assessed, but some differences between EBT and
25
26 eNO were observed: compared to the healthy controls, EBT was increased in all asthmatic
27
28 subjects, while eNO was only increased in those patients on inhaled corticosteroids,
29
30 suggesting that these two methods are picking different subtypes of asthmatic inflammation.
31
32
33

‘Turning EBT down’: the reduced airways heating surface

34
35
36 Histological studies of the airways in COPD patients found decreased bronchial
37
38 vascularity, the major determinant of EBT: reduction of the number of capillaries surrounding
39
40 the alveoli, structural changes in the small pulmonary arteries comprising the hypertrophy of
41
42 the inner membrane of secondary arteries and the smooth muscle cells [38]. These changes
43
44 result from repeated bouts of inflammation and lead to airway remodeling. The proposed
45
46 sequence of events involves signals from the damaged airway epithelium which elicit
47
48 immunological responses mobilizing underlying mesenchymal cells to start tissue repair. As
49
50 different types of inflammation (driven by specific sensitization to allergens, environmental
51
52 hazards or infection) persist or frequently recur in chronic respiratory diseases, the repair
53
54 process turns pathological in the course of time [39]. The repair process itself is not strictly
55
56 defined and may exhibit specific features along the continuum from the upper airways to the
57
58
59
60

1
2
3 respiratory bronchioles in the periphery of the bronchial tree. This may also be influenced by
4
5 the airway geometry and by structural differences in the extracellular matrix (ECM) scaffold
6
7 along the cascade of branching airways. This hypothesis was confirmed by Churg et al., who
8
9 suggested that genes involved in tissue repair were up-regulated in small airways but were
10
11 differentially expressed or down-regulated in the lung parenchyma after exposure to cigarette
12
13 smoke [40]. Furthermore, some studies have identified phenotypically unique subpopulations
14
15 of fibroblasts, key players in tissue repair, in central airways and in the parenchyma [41, 42].
16
17 The end result, the imperfect repair, is tissue remodeling with loss of elastic recoil,
18
19 degradation of alveolar walls (i.e. emphysema) and substantial heterogeneity of lung function
20
21 and gas volumes with gas trapping, further potentiated by hypoxia [43]. All these pathological
22
23 mechanisms impair the thermal exchange between the airway wall and the flow of air and
24
25 decrease more or less EBT. Contributors to this decrease are the thickened basement
26
27 membrane, the reduced vascular bed and eventually the hyperproduction of mucus [44, 45].
28
29 The relatively low EBT when the disease is under control may surge again when a new
30
31 inflammatory episode occurs.
32
33
34
35
36
37

EBT in respiratory diseases

38
39
40
41 The usefulness of single time-point measurement for diagnostic purposes is limited by
42
43 the mere nature of the processes shaping EBT: “pure” airway inflammation on the one hand
44
45 and reduction of the thermal convection airway surface (remodeling, tissue destruction, excess
46
47 mucus, shutting out of lung segments by obstruction/plugging) on the other, are two extremes
48
49 with a broad gray area between them in different lung disease entities [47] (Figure 3):
50
51
52
53
54
55
56
57
58
59
60

Exhaled breath temperature in humans




Figure 2. EBT ranges in healthy subjects and in patients with different respiratory diseases.

Asthma

Most studies on EBT so far have been done in asthmatic patients and have sought association with different anthropometric, clinical, lung function, laboratory and quality of life indices (Table 2).

Association of EBT with:		References
Age:		
- children	Yes	18, 50
- adults (19-60 years)	No	11,14
- elderly (>60 years)	Yes	16
Symptoms	Yes	8, 9, 11, 47, 48, 49, 54, 55, 56, 57, 58, 59
Spirometry/ FEV1	No	8, 9, 10, 47, 48, 49, 54, 55, 56, 57, 58, 59
FeNO	Yes	8, 9, 37, 48
Sputum eosinophils	Yes	9, 48, 56
Blood:		
- eosinophils	Yes	55, 57
- CRP	Yes	55, 57
- periostin	Yes	55

Figure 2. Association of EBT with basic patient characteristics in patients with asthma.

All studies have found increased EBT in asthmatic patients compared to control subjects without respiratory diseases in both adults and children, thus substantiating the utility of this approach to non-invasively assess airway inflammation.

Exhaled breath temperature in humans

1
2
3 Garcia et al. documented significantly higher EBT in 50 patients with uncontrolled
4 asthma compared with 50 patients with controlled asthma, EBT in both of these groups being
5 significantly higher than in 50 healthy controls [47].
6
7
8

9
10 Piacentini et al. [48] and Leornardi et al. [49] found significantly higher EBT in
11 asthmatic children and matched healthy controls. A significant positive relationship was also
12 observed by the Piacentini team between EBT and both exhaled nitric oxide and the
13 percentage of eosinophils in samples from induced sputum [48].
14
15
16
17
18

19
20 However, single point EBT measurements did not differ significantly in 134 asthmatic
21 children in terms of asthma control and treatment decisions by their physicians [50, Hamill].
22 A possible reason for this negative finding could be the confounding role that age plays in
23 childhood [18, 51].
24
25
26
27
28

29
30 Piacentini et al. were the first to come up with the hypothesis about a relationship
31 between EBT and airway remodeling in children/adolescence. In one study they found a
32 significant correlation between EBT and metalloproteinase-9 in asthmatic children [52], and
33 in another trial they documented a significant negative correlation between EBT and
34 diffusion lung capacity of carbon monoxide (DL_{CO}) [53].
35
36
37
38
39
40

41
42 Introducing repeated measurements over time provides added value to the EBT
43 method. Comparing EBT in patients with EBT measurement reveals asthma improvement in
44 the course of the anti-inflammatory treatment. This has been documented for pharmaceutical
45 products [11,54,55], for specific allergen immunotherapy [56] and for an acoustic medical
46 device mobilizing secretions from the lower airways of asthmatics [57]. Similarly to serial
47 peak expiratory flow (PEF) measurement, it follows a day-to-day pattern in line with the
48 control of asthma [58].
49
50
51
52
53
54
55
56

57
58 In a recent study EBT has shown promise as a marker and predictor of asthma
59 exacerbation in children and adolescents [59]. However, studies on natural exacerbations
60

Exhaled breath temperature in humans

1
2
3 require serial measurements and are rather difficult to design and implement due to the
4
5 unpredictability of these events. Alternatively, studies withholding asthma medications to
6
7 precipitate mild exacerbations of asthma face ethical issues. The availability of affordable
8
9 patient-friendly devices for daily EBT monitoring would reveal whether exacerbations could
10
11 be reliably predicted so as to provide a window of opportunity for early preventive measures.
12
13

14
15 Exercise, particularly in children, can elicit bronchoconstriction and is used in clinical
16
17 practice to prove the existence of airway hyperresponsiveness. Peroni et al. demonstrated that
18
19 EBT rises significantly after a standardized exercise test [60]. Two studies have addressed the
20
21 effect of exercise on EBT in asthmatic swimmers [61,62]. After a training session EBT
22
23 increased both in the athletes with and without asthma. However, in the study of Svenson et
24
25 al. EBT remained higher in the asthmatics whose FEV₁ dropped by >10% compared to the
26
27 remaining asthmatics [61].
28
29
30

COPD

31
32
33 The notion that EBT may also be affected by airway remodeling was supported by
34
35 data in patients with chronic degenerative respiratory disease. Paredi et al. were the first to
36
37 report slower rise of exhaled breath temperature in COPD [63]. A study of adolescents who
38
39 survived bronchopulmonary dysplasia found their EBT to be significantly lower than in age
40
41 matched asthmatics, suggesting that different pathogenetic mechanisms characterize this
42
43 chronic obstructive disease state [64]. Kløkstad et al. found significantly lower EBT in COPD
44
45 patients compared with smokers and healthy controls, which made them suggest that even
46
47 though airway inflammation was present in this disease, the structural damage of
48
49 airway/alveolar tissue with consequently impaired blood flow might have resulted in an
50
51 overall lower breath temperature [65]. This notion was further illustrated by the same team by
52
53 demonstrating that when COPD patients exacerbated, this still led to an increase of EBT [66].
54
55
56
57
58
59
60 The same pattern was found by Labor et al. [28] but the difference between groups ('healthy'

1
2
3 smokers, symptomatic smokers and COPD GOLD stage-I did not reach statistical
4
5 significance. The prolonged duration of the measurement procedure with the multiple breath
6
7 EBT measurement device provides indirect evidence of the lower potential of the airways to
8
9 “heat up” the outgoing air [67].
10
11

12 Lázár Z et al. found distinct differences in EBT of patients with stable COPD, of
13
14 patients with COPD at onset and also after recovery from an acute exacerbation, of control
15
16 smoking/ex-smoking control subjects [69]. Patients with stable COPD had lower EBT values
17
18 than smokers/ex-smokers. EBT was higher at the onset of acute exacerbations of COPD
19
20 compared to the patients in a stable condition, and decreased after recovery. The increased
21
22 EBT during exacerbations positively correlated with the sputum leukocyte counts.
23
24
25
26

27 Labor et al. were first to investigate EBT as a susceptibility marker to cigarette smoke
28
29 in order to predict COPD development in smokers at risk [28]. Results of this study showed
30
31 the potential of a change in EBT from baseline, after smoking a cigarette (Δ EBT), to be
32
33 significantly predictive for development of manifest COPD and for the disease progression
34
35 after 2 years. The same team is extending their EBT studies in an attempt to establish the
36
37 COPD diagnosis in the pre-symptomatic stage, before significant end organ damage [70].
38
39
40

41 **Cystic fibrosis**

42
43 Cystic fibrosis (CF) is characterized by chronic airway infection and inflammation
44
45 pushing EBT up, and structural changes of the airways and lung tissues, pushing it down.
46
47 Subsequently, Garcia et al. did not find significant differences between the EBT of adult CF
48
49 patients and healthy controls [71]. Similarly, Bade et al. did not find differences in the
50
51 absolute EBT values of patients and controls, but established a slower rise of EBT in CF
52
53 patients [72].
54
55
56

57
58 A multinational team studied 57 CF patients and measured their EBT by a single-
59
60 breath method. They also assessed the temperature of sputum and directly of the airway

Exhaled breath temperature in humans

1
2
3 lumen and wall using fiberoptic bronchoscopy [73]. The investigators found a significant
4
5 inverse correlation between EBT and FEV₁, with EBT values of the more obstructed subjects
6
7 higher than those of the controls.
8
9

Lung cancer

10
11
12 Cancerous growth in the lungs is characterized by inflammation and increased
13
14 vascularity. Carpagnano et al. measured EBT in 82 consecutive patients with suspected non-
15
16 small-cell lung cancer (NSCLC) in order to explore the applicability of the method for
17
18 diagnostic and monitoring purposes [74]. In 40 patients cancer diagnosis was confirmed by
19
20 the standard work-up, while the remaining 42 were labeled as false-positive and were used as
21
22 controls. EBT turned out to be significantly higher in the NSCLC patients compared to the
23
24 healthy subjects. EBT was correlated with the number of pack-years and associated with the
25
26 stage of lung cancer. The authors determined the cut-off value for EBT that could screen
27
28 patients with lung cancer with high sensitivity and specificity.
29
30
31
32

33
34 In a subsequent study the same team enrolled 44 consecutive patients with
35
36 radiological suspicion of lung cancer and ten healthy non-smoker volunteers, in all of whom
37
38 EBT was measured [75]. The researchers also measured leukotriene B₄, a marker of airways
39
40 inflammation, and vascular endothelial growth factor (VEGF), a marker of neoangiogenesis,
41
42 in exhaled breath condensate. They confirmed the previous finding of a higher EBT in lung
43
44 cancer patients compared to the controls. A multiple linear regression model showed that the
45
46 exhaled VEGF was the only predictor of elevated of EBT, which they interpreted as evidence
47
48 that angiogenesis was driving the increase in EBT in lung cancer. The study suggested the
49
50 potential for use of EBT in monitoring lung cancer progression.
51
52
53

Infections

54
55
56 Both viral and bacterial infections have a direct bearing on exacerbations of chronic
57
58 lung diseases. Infections can be confined to the respiratory system, but can also be associated
59
60

Exhaled breath temperature in humans

1
2
3 with systemic symptoms including general febrile episodes, so it is important to be aware of
4 the relationship between fever and EBT. Six generally healthy subjects measured their EBT
5 and ear temperature (ET) daily for periods of between 5 months and 2 years, using personal
6 hand-held devices uploading the results on a specialized web site [76]. They were instructed
7 to start recording both ET and EBT at 8-hour intervals if they felt signs of a general
8 indisposition and if their ET exceeded 37°C. Six episodes of fever were documented during
9 the study: 2 cases of rhinovirus infections in which EBT rose by 1.2-1.9°C above baseline,
10 preceding by 24-72 hours a moderate increase of ET of up to 38°C; 2 cases of influenza in
11 which EBT rose by >2.0°C about 6 hours before increase of ET up to 40°C; 2 cases of
12 bacterial infections, urinary and GI, during which EBT rose by ≈1.0°C simultaneously with
13 the rise of ET (up to 39°C). These results prompted the conclusion that EBT rises during viral
14 infections, affecting the respiratory system earlier than ET, providing a window of
15 opportunity for early treatment. This may have implications for patients at risk of
16 exacerbations of underlying obstructive airway diseases. The method may also discriminate
17 between different disease agents, which warrant specific research designs.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
The issue of asthma exacerbations in the pediatric range was explored by Xepapadaki
et al. [77]. They documented significant EBT increase at the onset of virus triggered asthma
exacerbations. The possibility of using personal devices for EBT measurement opens the door
for prospective studies to assess the value of serial home measurements. EBT measurements
may predict in advance the onset of viral infections, providing the opportunity to prevent or
abate subsequent exacerbations.

Applicability of EBT measurement

The idea that motivated initial research into the measurement of EBT was rather
simple and straightforward: as airway inflammation has gained unanimous recognition as the
hallmark of asthma, and as increased temperature is a prominent feature of inflammation,

Exhaled breath temperature in humans

1
2
3 detecting the thermal signal from the inflamed airways would be a simple measure of the
4
5 state of asthma control. Insight into the nature of the processes shaping EBT gained
6
7 complexity as data started to accumulate over time [46]. An important element configuring
8
9 the EBT model is airway remodeling. This is in contrast to FeNO, the closest approximation
10
11 of what EBT measurement can be used for, which is associated exclusively with eosinophilic
12
13 airway inflammation and hyperresponsiveness. In fact, in cases of advanced chronic lung
14
15 disease, where FeNO has little value, EBT can get quite low, thus adding an important
16
17 dimension to the applicability of the method. As a matter of fact, these two non-invasive
18
19 methods can be used conjointly to detect the eosinophilic airway inflammation signal by
20
21 assessing increased FeNO in subjects with decreased EBT due to airway remodeling.
22
23 Baseline EBT is compounded by the processes of inflammation and remodeling, which act in
24
25 opposite directions and sometimes the resulting vector could be lying within the “normal”
26
27 range. From this point of view the broad clinical spectrum of chronic airway diseases should
28
29 be regarded as individual combinations of inflammation and remodeling. Documenting EBT
30
31 at a time point of adequate disease control may serve as a reference point to warn of
32
33 imminent inflammatory exacerbation, and of advancement of remodeling/destruction in the
34
35 long run. A prerequisite to this end would be monitoring with user friendly individual devices
36
37 for EBT measurement.
38
39
40
41
42
43
44
45

Future perspectives

46
47
48
49 EBT, which was initially an unexplored area on the map of human physiology and
50
51 disease, is gradually being completed. It has the potential to satisfy the need of current clinical
52
53 practice in the field of lung diseases for a simple, cheap and non-invasive tool to assess and
54
55 monitor the state of the airways. Ongoing systematic research will determine its place in the
56
57 clinical setting and as a tool in home monitoring in line with the modern trends of
58
59 personalized and telemedicine. The usefulness of this approach should be enhanced by either
60

1
2
3 combining it with other objective measurements, or by breaking it down to components that
4
5 could possibly differentiate between phenotypes of airway diseases. This second option
6
7 prompted the idea of assessing separately the contribution of the central and peripheral
8
9 airways to compliment the standard integral EBT measurement [78]. This will help avoid
10
11 false-negative findings of 'normal' EBT in subjects with equal part of both inflammatory and
12
13 degenerative disorders. It will also help evaluate the kinetics of different diagnostic and
14
15 therapeutic approaches, adding a new dimension in differentiating health from airway
16
17 pathology.
18
19
20
21

22 23 **References**

- 24
25 1. Wilson AP. A treatise on Febrile Diseases, including intermitting, remitting, and continued
26 fevers; eruptive fevers; inflammations; hemorrhagies; and the piofluvia; in which an attempt is
27 made to present, at one view, whatever, in the present state of medicine, it is requisite for
28 physician to know respecting the symptoms, causes, and cure of those diseases. 1800,
29 Monograph, Printed and sold by Robbins, London, Edinburgh.
- 30
31 2. Sund-Levander M, Grodzinsky E. Time for a change to assess and evaluate body temperature in
32 clinical practice. *Int J Nurs Pract.* 2009; 15(4): 241-9.
- 33
34 3. Tansey EA, Johnson CD. Recent advances in thermoregulation. *Adv Physiol Educ.* 2015; 39
35 (3): 139-48.
- 36
37 4. Sund-Levander M, Forsberg C, Wahren LK. Normal oral, rectal, tympanic and axillary body
38 temperature in adult men and women: a systematic literature review. *Scand J Caring Sci.* 2002;
39 16 (2): 122-8.
- 40
41 5. Mallory M, Gogineni E, Jones GC, Greer L, Simone CB 2nd. Therapeutic hyperthermia: The
42 old, the new, and the upcoming. *Crit Rev Oncol Hematol.* 2016; 97: 56-64.
- 43
44 6. Insler SR, Sessler DI. Perioperative thermoregulation and temperature monitoring. *Anesthesiol*
45 *Clin.* 2006; 24(4): 823-37.
- 46
47 7. Barnett BJ, Nunberg S, Tai J, Lesser ML, Fridman V, Nichols P, Powell R, Silverman R. Oral
48 and tympanic membrane temperatures are inaccurate to identify fever in emergency department
49 adults. *West J Emerg Med.* 2011; 12 (4): 505-11.
- 50
51 8. Paredi P, Kharitonov SA, Barnes PJ. Faster rise of EBT in asthma: a novel marker of airway
52 inflammation? *Am J Respir Crit Care Med* 2002; 165: 181–184.
- 53
54 9. Piacentini GL, Bodini A, Zerman L, Costella S, Zanolla L, Peroni DG, Boner AL.. Relationship
55 between exhaled air temperature and exhaled nitric oxide in childhood asthma. *Eur Respir J*
56 2002; 20: 108–111.
- 57
58 10. Pifferi M, Ragazzo V, Previti A, Pioggia G, Ferro M, Macchia P, Piacentini GL, Boner AL.
59 Exhaled air temperature in asthmatic children: a mathematical evaluation. *Pediatr Allergy*
60 *Immunol.* 2009; 20 (2): 164-71.
11. Popov TA, Dunev SS, Kralimarkova TK, Kraeva S, DuBuske LM. Evaluation of a simple,
potentially individual device for exhaled breath temperature measurement. *Respiratory*
Medicine. 2007; 101 (10): 2044-2050.

Exhaled breath temperature in humans

12. Popov TA, Kralimarkova TZ, Tzachev CT, Dimitrov VD, Mun KK, Gill J. Exhaled breath temperature measurement made easy. *Pediatr Allergy Immunol.* 2009; 20 (2): 200-201.
13. Popov TA, Kralimarkova TZ, Tzachev CT, Dunev S, Dimitrov,VD, Gill J. Development of an individual device for exhaled breath temperature measurement. *IEEE Sensors Journal*, 2010; 10 (1): 110-3.
14. Carpagnano GE, Foschino-Barbaro MP, Crocetta C, Lacedonia D, Saliani V, Zoppo LD, Barnes PJ. Validation of the exhaled breath temperature measure: reference values in healthy subjects. *Chest.* 2016 Nov 23.
15. Logie KM, Kusel MMH, Sly PD, Hall GL. Exhaled breath temperature in healthy children is influenced by room temperature and lung volume. *Pediatr Pulmonol*, 2011; 46 (11): 1062–1068.
16. Bijnens E, Pieters N, Dewitte H, Cox B, Janssen BG, Saenen N, Dons E, Zeegers MP, Int Panis L, Nawrot TS. Host and environmental predictors of exhaled breath temperature in the elderly. *BMC Public Health.* 2013; 13: 1226.
17. Flouris AD, Cheung SS. The validity of tympanic and exhaled breath temperatures for core temperature measurement. *Physiol Meas.* 2010; 31(5): N35-42.
18. Popov TA, Kralimarkova TZ. Exhaled breath temperature measurement: applicability in childhood. *Pediatr Pulmonol.* 2016. 51(1): 91-2. 38.
19. Barreto M, Piacentini G, Chioffi L, Ruggeri F, Caiazzo I, Campisano M, Martella S, Villa MP. Tidal-breathing measurement of exhaled breath temperature (EBT) in schoolchildren. *Pediatric Pulmonology.* 2014; 49(12): 1196-204.
20. Vermeulen S, Barreto M, La Penna F, Prete A, Martella S, Biagiarelli F, Villa MP. Exhaled breath temperature in children: reproducibility and influencing factors. *J Asthma.* 2014; 51(7): 743-50.
21. Crespo Lessmann A, Giner J, Torrego A, Mateus E, Torrejón M, Belda A, Plaza V. Usefulness of the exhaled breath temperature plateau in asthma patients. *Respiration.* 2015; 90 (2): 111-7.
22. Kralimarkova TZ, Rasheva M, Grigorova T, Dimitrov Z, Tihomirov D, Mincheva R, Dimitrov VD, Popov TA. Circadian variation of exhaled breath temperature in healthy subjects. *ERS Congress 2011, Amsterdam, The Netherlands; 27 September 2011: 736s.*
23. Kralimarkova T, Mincheva R, Kadavil R, John J, Tihomirov D, Dimitrov V, Popov TA. Effect of energy food intake on exhaled breath temperature in healthy subjects. *Eur Respir J* 2012; 40: Suppl. 56, 631s.
24. Tufvesson E, Svensson H, Ankerst J, Bjermer L. Increase of club cell (Clara) protein (CC16) in plasma and urine after exercise challenge in asthmatics and healthy controls, and correlations to exhaled breath temperature and exhaled nitric oxide. *Respir Med.* 2013; 107(11): 1675-81.
25. Kralimarkova TZ, Mincheva RK, Dimitrov VD, Gill J, Popov T. Exhaled breath temperature and tobacco smoking. *Am J Respir Crit Care Med.* 2010; 181: A5439.
26. Juric I, Labor M, Labor S, Plavec D. Dynamics of exhaled breath temperature (EBT) after a smoked cigarette. *Eur Respir J.* 2016;48 (Suppl 60): OA3502.
27. Carpagnano GE, Ruggieri C, Scioscia G, Storto MM, Zoppo L, Foschino-Barbaro MP. Is the exhaled breath temperature sensitive to cigarette smoking? *COPD.* 2016; 13 (5): 642-6.
28. Labor M, Vrbica Ž, Gudelj I, Labor S, Jurić I, Plavec D. Exhaled breath temperature as a novel marker of future development of COPD: results of a follow-up study in smokers. *COPD.* 2016; 13 (6): 741-9.
29. Kralimarkova T, Dimitrov V, Popov T. Changes in exhaled breath temperature before, during and after the pollen season in subjects sensitised to grasses with rhinoconjunctivitis, with or without asthma. *Allergy* 2011; 66: 354s.

- 1
- 2
- 3 30. Kralimarkova TZ, Garcia G, Grigorova T, Yañez A, Bergna M, Mincheva R, Dimitrov VD,
- 4 Popov TA. Effect of an inhaled short acting beta-agonist on exhaled breath temperature. ERS
- 5 Congress 2010, Barcelona, Spain; 19 September 2010: 230s.
- 6
- 7 31. Svensson H, Nilsson D, Bjermer L, Tufvesson E. Exhaled breath temperature increases after
- 8 exercise in asthmatics and controls. *Respiration*. 2012; 84(4): 283-90.
- 9
- 10 32. Vignola AM, Mirabella F, Costanzo G, Di Giorgi R, Gjomarkaj M, Bellia V, Bonsignore G.
- 11 Airway remodeling in asthma. *Chest*. 2003; 123(3 Suppl): 417S-22S.
- 12
- 13 33. Salvato G. Quantitative and morphological analysis of the vascular bed in bronchial biopsy
- 14 specimens from asthmatic and non-asthmatic subjects. *Thorax*. 2001; 56(12):902-6.
- 15
- 16 34. Parsons GH, Nichol GM, Barnes PJ, Chung KF. Peptide mediator effects on bronchial blood
- 17 velocity and lung resistance in conscious sheep. *J Appl Physiol* (1985). 1992; 72(3):1118-22.
- 18
- 19 35. Charan NB, Johnson SR, Lakshminarayan S, Thompson WH, Carvalho P. Nitric oxide and
- 20 beta-adrenergic agonist-induced bronchial arterial vasodilation. *J Appl Physiol* (1985). 1997;
- 21 82(2): 686-92.
- 22
- 23 36. Tansey EA, Johnson CD. Recent advances in thermoregulation. *Adv Physiol Educ*. 2015; 39
- 24 (3): 139-48.
- 25
- 26 37. Paredi P, Kharitonov SA, Barnes PJ. Correlation of exhaled breath temperature with bronchial
- 27 blood flow in asthma. *Respiratory Research* 2005; 6, 15: 1-10.
- 28
- 29 38. Araya J, Nishimura SL. Fibrogenic reactions in lung disease. *Annu Rev Pathol*. 2010; 5: 77-98.
- 30
- 31 39. Fahy JV, Corry DB, Boushey HA. Airway inflammation and remodeling in asthma. *Curr Opin*
- 32 *Pulm Med*. 2000; 6(1): 15-20.
- 33
- 34 40. Chung A, Zhou S, Preobrazhenska O, Tai H, Wang R, Wright JL. Expression of profibrotic
- 35 mediators in small airways versus parenchyma after cigarette smoke exposure. *Am J Respir Cell*
- 36 *Mol Biol*. 2009; 40(3): 268-76.
- 37
- 38 41. Pechkovsky DV, Hackett TL, An SS, Shaheen F, Murray LA, Knight DA. Human lung
- 39 parenchyma but not proximal bronchi produces fibroblasts with enhanced TGF-beta signaling
- 40 and alpha-SMA expression. *Am J Respir Cell Mol Biol*. 2010; 43(6): 641-51.
- 41
- 42 42. Kotaru C, Schoonover KJ, Trudeau JB, Huynh ML, Zhou X, Hu H, Wenzel SE. Regional
- 43 fibroblast heterogeneity in the lung: implications for remodeling. *Am J Respir Crit Care Med*.
- 44 2006; 173 (11): 1208-15.
- 45
- 46 43. Polosukhin VV, Lawson WE, Milstone AP, Egunova SM, Kulipanov AG, Tchuvakin SG,
- 47 Massion PP, Blackwell TS. Association of progressive structural changes in the bronchial
- 48 epithelium with subepithelial fibrous remodeling: a potential role for hypoxia. *Virchows Arch*.
- 49 2007; 451(4): 793-803.
- 50
- 51 44. Peinado VI, Pizarro S, Barbera JA. Pulmonary Vascular Involvement in COPD. *Chest* 2008:
- 52 134: 808-14.
- 53
- 54 45. Allinson JP, Hardy R, Donaldson GC, Shaheen SO, Kuh D, Wedzicha JA. The Presence of
- 55 Chronic Mucus Hypersecretion across Adult Life in Relation to Chronic Obstructive Pulmonary
- 56 Disease Development. *Am J Respir Crit Care Med*. 2016; 193(6): 662-72.
- 57
- 58 46. Kralimarkova TZ, Popov TA. Exhaled breath temperature: broadening the horizons. *Int J*
- 59 *Tuberc Lung Dis*, 2014; 18(2): 250-1.
- 60
47. García G, Bergna M, Uribe E, Yañez A, Soriano JB. Increased exhaled breath temperature in
- subjects with uncontrolled asthma. *Int J Tuberc Lung Dis*. 2013; 17(7): 969-72.
48. Piacentini GL, Peroni D, Crestani E, Zardini F, Bodini A, Costella S, Boner AL. Exhaled air
- temperature in asthma: methods and relationship with markers of disease. *Clin Exp Allergy*.
- 2007; 37(3):415-9.

Exhaled breath temperature in humans

- 1
2
3 49. Leonardi S, Cuppari C, Lanzafame A, Attardo D, Tardino L, Parisi G, Giacchi V, Manti S,
4 Arrigo T. Exhaled breath temperature in asthmatic children. *J Biol Regul Homeost Agents*.
5 2015; 29 (2 Suppl 1): 47-54.
6
- 7 50. Hamill L, Ferris K, Kapande K, McConaghy L, Douglas I, McGovern V, Shields MD. Exhaled
8 breath temperature measurement and asthma control in children prescribed inhaled
9 corticosteroids: A cross sectional study. *Pediatr Pulmonol*. 2016; 51(1): 13-21.
10
- 11 51. Hamill L, Ferris K, Kapande K, McConaghy L, Douglas I, McGovern V, Shields MD. Response
12 to letter by Popov, Todor regarding our paper: Exhaled breath temperature measurement and
13 asthma control in children prescribed inhaled corticosteroids: A cross sectional study. *Pediatr*
14 *Pulmonol*. 2016; 51(1): 93.
15
- 16 52. Piacentini GL, Peroni DG, Bodini A, Corradi M, Boner AL. Exhaled breath temperature as a
17 marker of airway remodelling in asthma: a preliminary study. *Allergy*. 2008; 63(4): 484-5.
18
- 19 53. Piacentini GL, Tezza G, Cattazzo E, Kantar A, Ragazzo V, Boner AL, Peroni DG. Diffusion
20 lung capacity of carbon monoxide: A novel marker of airways remodeling in asthmatic
21 children? *Allergy Rhinol (Providence)*. 2012; 3(2): e66-73.
22
- 23 54. Melo RE, Popov TA, Solé D. Exhaled breath temperature, a new biomarker in asthma control: a
24 pilot study. *J Bras Pneumol*. 2010; 36(6): 693-9.
25
- 26 55. Popov TA, Petrova D, Kralimarkova TZ, Ivanov Y, Popova T, Peneva M, Odzhakova T, Ilieva
27 Y, Yakovliev P, Lazarova T, Georgiev O, Hodzhev V, Hodzheva E, Staevska MT, Dimitrov VD.
28 Real life clinical study design supporting the effectiveness of extra-fine inhaled
29 beclomethasone/formoterol at the level of small airways of asthmatics. *Pulm Pharm*
30 *Therapeutics* 2013; 26: 624-29.
31
- 32 56. Kralimarkova TZ, Popov TA, Staevska M, Mincheva R, Lazarova C, Racheva R, Mustakov TB,
33 Filipova V, Koleva M, Bacheva K, Dimitrov VD. Objective approach for fending off the
34 sublingual immunotherapy placebo effect in subjects with pollenosis: double-blinded, placebo-
35 controlled trial. *Ann Allergy Asthma Immunol* 2014; 113 (1): 108-13.
36
- 37 57. Kralimarkova TZ, Dimitrov Z, Koleva M, Filipova V, Rasheva M, Gugutkova M, Mincheva R,
38 Dimitrov VD, DuBuske LM, Popov TA. Increase of exhaled breath temperature after
39 mobilizing secretions from the lower airways of asthmatics using acoustic wave technology.
40 *Ann Allergy Asthma Immunol*. 2012; 109 (5): A59.
41
- 42 58. Popov TA, Kralimarkova TZ, Lazarova C, Tzachev CT, Dimitrov,VD. Daily monitoring of
43 asthmatics by means of individual devices for exhaled breath. *IEEE Sensors Journal*, 2010; 10
44 (1): 44-8.
45
- 46 59. Wojsyk-Banaszak I, Mikoś M, Szczepankiewicz A, Wielebska A, Sobkowiak P, Kamińska A,
47 Bręborowicz A. Evaluation of exhaled breath temperature (EBT) as a marker and predictor of
48 asthma exacerbation in children and adolescents. *J Asthma*. 2017; 10: 1-7.
49
- 50 60. Peroni DG, Chinellato I, Piazza M, Zardini F, Bodini A, Olivieri F, Boner AL, Piacentini GL.
51 Exhaled breath temperature and exercise-induced bronchoconstriction in asthmatic children.
52 *Pediatr Pulmonol*. 2012; 47 (3): 240-4.
53
- 54 61. Svensson H, Nilsson D, Bjermer L, Tufvesson E. Exhaled breath temperature increases after
55 exercise in asthmatics and controls. *Respiration*. 2012; 84(4): 283-90.
56
- 57 62. Svensson H, Bjermer L, Tufvesson E. Exhaled breath temperature in asthmatics and controls
58 after eucapnic voluntary hyperventilation and a methacholine challenge test. *Respiration*. 2014;
59 87(2): 149-57.
60
63. Couto M, Santos P, Silva D, Delgado L, Moreira A. Exhaled breath temperature in elite
swimmers: the effects of a training session in adolescents with or without asthma. *Pediatr*
Allergy Immunol. 2015; 26(6): 564-70.

- 1
2
3 64. Paredi P, Caramori G, Cramer D, Ward S, Ciaccia A, Papi A, Kharitonov SA, Barnes PJ.
4 Slower rise of exhaled breath temperature in chronic obstructive pulmonary disease. *Eur Respir*
5 *J.* 2003; 21(3): 439-43.
6
7 65. Carraro S, Piacentini G, Lusiani M, Uyan ZS, Filippone M, Schiavon M, Boner AL, Baraldi E.
8 Exhaled air temperature in children with bronchopulmonary dysplasia. *Pediatr Pulmonol.* 2010;
9 45(12): 1240-5.
10
11 66. Kløkstad S, Bikov A, Lazar Z, Galffy G, Losonczy G, Horvath I. The effect of smoking and
12 COPD on exhaled breath temperature. *ERS Congress 2010, Barcelona, Spain; 21 September*
13 *2010: 832s.*
14
15 67. Lázár Z, Bikov A, Gálffy G, Orosz M, Losonczy G, Hováth I. Exhaled breath temperature
16 increases at COPD exacerbation and correlates with sputum neutrophilia. *ERS Congress 2011,*
17 *Amsterdam, The Netherlands; 27 September 2011: 874-5s.*
18
19 68. Kralimarkova TZ, Tzachev CT, Dimitrov VD, Popov TA. Duration of exhaled breath
20 temperature measurement with a hand-held device as a physiological index on its own. *Eur*
21 *Respir J.* 2009; 34, Suppl. 53: 564s-566s.
22
23 69. Lázár Z, Bikov A, Martinovszky F, Gálffy G, Losonczy G, Horváth I. Exhaled breath
24 temperature in patients with stable and exacerbated COPD. *J Breath Res.* 2014; 8(4): 046002.
25
26 70. Vrbica Ž, Labor M, Gudelj I, Labor S, Jurić I, Plavec D; MARKO study group. Early detection
27 of COPD patients in GOLD 0 population: an observational non-interventional cohort study -
28 MARKO study. *BMC Pulm Med.* 2017; 17 (1): 36.
29
30 71. Garcia G, Granero N, Hendriksen B, Carlos D, Ezequiel B, Bergna M. Exhaled breath
31 temperature in adult cystic fibrosis. *ERS Congress 2011, Amsterdam, The Netherlands; 27*
32 *September 2011: 788s.*
33
34 72. Bade G, Gupta S, Kabra SK, Talwar A. Slower rise of exhaled breath temperature in cystic
35 fibrosis. *Indian Pediatr.* 2015; 52(2): 125-7.
36
37 73. Schmidt A, Belaaouaj A, Bissinger R, Koller G, Malleret L, D'Orazio C, Facchinelli M,
38 Schulte-Hubbert B, Molinaro A, Holst O, Hammermann J, Schniederjans M, Meyer KC,
39 Damkiaer S, Piacentini G, Assael B, Bruce K, Häußler S, LiPuma JJ, Seelig J, Worlitzsch D,
40 Döring G. Neutrophil elastase-mediated increase in airway temperature during inflammation. *J*
41 *Cyst Fibros.* 2014; 13 (6): 623-31.
42
43 74. Carpagnano GE, Lacedonia D, Spanevello A, Martinelli D, Saliari V, Ruggieri C, Foschino-
44 Barbaro MP. Exhaled breath temperature in NSCLC: could be a new non-invasive marker?
45 *Med Oncol.* 2014; 31(5): 952.
46
47 75. Carpagnano GE, Lacedonia D, Spanevello A, Cotugno G, Saliari V, Martinelli D, Foschino-
48 Barbaro MP. Is the exhaled breath temperature in lung cancer influenced by airways
49 neoangiogenesis or by inflammation? *Med Oncol.* 2015; 32(10): 237.
50
51 76. Popov TA, Kralimarkova TZ, DuBuske LM. Relationship between exhaled breath temperature
52 and ear temperature in otherwise healthy persons during febrile infectious illness. *J Allergy Clin*
53 *Immunology* 2016; 137 (2, Suppl.): AB202.
54
55 77. Xepapadaki P, Xatziioannou A, Chatzicharalambous M, Makrinioti H, Papadopoulos NG.
56 Exhaled breath temperature increases during mild exacerbations in children with virus-induced
57 asthma. *Int Arch Allergy Immunol.* 2010; 153(1): 70-4.
58
59 78. Popov TA, Kralimarkova TZ, Hristova D, et al. Single breath method to assess the relative
60 contribution of central and peripheral airways in the overall exhaled breath temperature. *J*
Allergy Clin Immunology 2015; 135 (2, Suppl.): AB177.